

An analysis of the significant variation in psychostimulant use across the U.S.[†]

Farasat Bokhari PhD¹, Rick Mayes PhD^{2*} and Richard M. Scheffler PhD³

¹*Department of Economics, Florida State University, FL, USA*

²*Department of Political Science, University of Richmond, Richmond, VA, USA*

³*Department of Economics & Public Policy, Graduate School of Public Health, University of California, Berkeley, CA, USA*

SUMMARY

Objective To provide a national profile of the area variation in per-capita psychostimulant consumption in the U.S.

Methods We separated 3030 U.S. counties into two categories of 'low' and 'high' per-capita use of attention deficit hyperactivity disorder (ADHD) drugs (based on data from the Drug Enforcement Administration), and then analyzed them on the basis of their socio-demographic, economic, educational and medical characteristics.

Results We found significant differences and similarities in the profile of counties in the U.S. that are above and below the national median rate of per-capita psychostimulant use (defined as g/per 100K population). Compared to counties below the median level, counties above the median level have: significantly greater population, higher per-capita income, lower unemployment rates, greater HMO penetration, more physicians per capita, a higher ratio of young-to-old physicians and a slightly higher students-to-teacher ratio.

Conclusions Our analysis of the DEA's ARCOS data shows that most of the significant variables correlated with 'higher' per-capita use of ADHD drugs serve as a proxy for county affluence. To provide a more complex, multivariate analysis of the area variation in psychostimulant use across the U.S.—which is the logical next step—requires obtaining price data to match the DEA's quantity data. Copyright © 2004 John Wiley & Sons, Ltd.

KEY WORDS — psychostimulants; attention deficit hyperactivity disorder (ADHD); methylphenidate; amphetamine; mental health; children; adolescents

BACKGROUND: PSYCHOSTIMULANT USE AS TREATMENT FOR ADHD

Psychostimulants (methylphenidate and amphetamines) are primarily used to treat attention deficit hyperactivity disorder (ADHD), which is the most commonly diagnosed behavioral disorder in children, making up more than 50% of all child psychiatric diagnoses.¹ Between 3 and 5% of U.S. school-age children

are estimated to have ADHD. But individual community studies have reported prevalence rates ranging from as low as 1.7 to as high as 26%.^{2,9,20–28} The core symptoms of ADHD include developmentally inappropriate levels of attention, concentration, activity, distractibility and impulsivity. According to the National Institutes of Health (NIH), children with ADHD have pronounced impairments and can experience long-term adverse effects on academic performance, vocational success and social-emotional development which have a profound impact on individuals, families, schools and society.³

ADHD is a major public health concern. Children with the disorder 'consume a disproportionate share of resources and attention from the health care system, criminal justice system, schools and other social

*Correspondence to: Dr Rick Mayes, PhD, Assistant Professor of Public Policy, 28 Westhampton Way, University of Richmond, Richmond, VA 23173, USA. E-mail: bmayes@richmond.edu

[†]No conflict of interest was declared.

Received 21 October 2003

Revised 24 March 2004

Accepted 5 May 2004




Figure 1. Methylphenidate distribution rate from selected states and national average: 1980–1996. (Source: Drug Enforcement Administration’s ARCOS database, 1980–1996)

service agencies’.³ The direct costs of medical care for children and adolescents with the disorder are substantial for their families. Children with ADHD incur double the amount of out-of-pocket expenses compared to children without the disorder, as well as averaging 10 times more outpatient mental health visits, 3.4 times more pharmacy fills and 1.6 times more primary care visits.⁴

Over the last decade and a half, psychostimulant use has increased dramatically (Figure 1),^{6,11} primarily

among children and adolescents but also increasingly among adults.¹² At the same time, there appears to be significant area variation in psychostimulant use (Table 1, Figure 2).²⁰ In addition to our own findings, consumption rates have previously been reported to vary as much as 1:4 between states and as high as 1:10 between communities within states.^{13,14}

The significant growth in psychostimulant use began in the early 1990s (Figure 1), soon after major changes were enacted by policymakers in Washington, D.C., to

Table 1. State psychostimulant distribution in kg/per 100 000 individuals, 2000

High quartile 75th percentile		Above average quartile 50–75th percentile		Below average quartile 25–50th percentile		Low quartile 25th percentile	
DE (7.4)	NH (7.1)	ME (5.1)	MD (5.1)	LA (4.5)	OH (4.4)	OK*(3.7)	CO (3.5)
RI*†(6.2)	VT (6.2)	IN* (5.1)	MO (5.0)	WA (4.4)	PA (4.3)	AR (3.3)	FL (3.3)
MA*(6.0)	IA (5.9)	NC (4.9)	GA (4.9)	ID*†(4.3)	TN (4.3)	DC (3.2)	SD (3.1)
MI (5.8)	SC (5.6)	AL (4.8)	KS (4.7)	WV*(4.3)	OR (4.2)	MS (3.0)	NJ (3.0)
WI (5.5)	MT (5.4)	ND (4.6)	UT*(4.6)	KY*(4.1)	TX*(4.1)	NM*(2.7)	CA*†(2.3)
VA (5.4)	MN (5.3)	AR (4.6)	CT (4.6)	NE (4.0)	IL*(3.9)	NV*(2.2)	NY*†(2.1)
		AK (4.6)		WY (3.9)		HI*†(1.7)	

Source: Drug Enforcement Administration’s ARCOS database, 2000.

*Designates the 15 states with a Schedule II triplicate prescription monitoring program.

†Designates the five states with monitoring programs in operation in 1981.

VARIATION IN PSYCHOSTIMULANT USE




Figure 2. Psychostimulant distribution rates 2000

include ADHD as a protected disability under the supplemental security income (SSI) program and the individuals with disabilities education act (IDEA). These changes in federal disability policy may have led to an increased awareness of the disorder among some teachers, parents and clinicians. They may have also decreased the stigma associated with the diagnosis. But between 1990 and 1996, psychostimulant consumption increased 370% nationwide,²⁹ while the number of patients diagnosed with the disorder grew from around 900 000 to approximately 3 million.³⁰

As Figure 1 illustrates, some states (Michigan, Georgia, Virginia) have had a tradition of using psychostimulants more than others (New York, California), while various states (Massachusetts, Texas and Pennsylvania) have oscillated above and below the national average from year to year. Viewed together with Figure 1, Table 1 shows the variation in distribution rates across states, which ranged from as high as 7.4 to as low as 1.7 kg/per 100 000 individuals in 2000.

Table 1 also shows that the 15 states with Schedule II triplicate prescription monitoring programs³¹ in 2000 had a combined psychostimulant distribution rate (3.3 kg/per 100 000 individuals) that was 36% lower than the combined distribution rate (4.5 kg/per 100 000 individuals) of the remaining 35 states, including D.C.,

which does not have a monitoring program. This statistically significant differential has existed every year, with minor fluctuations, from 1980 to 2000. Nevertheless, these same 15 states also used 36% less of the drugs back in 1981, when only five of them had monitoring programs in operation. So it is an open question as to how much, if at all, they dampen the aggregate use of psychostimulants.³² The programs could simply be the product of the states' traditional predilection for lower use of these drugs.

Ultimately, the purpose of our study is to understand more about how psychostimulant use varies by socio-economic, demographic, educational and health system characteristics. For instance, we want to know if psychostimulant use is higher in more affluent areas, as well as areas with more: clinicians, children as a proportion of the total population; students-per-teacher; more white children as a proportion of the overall childhood population and HMO penetration.

RESEARCH DESIGN AND DATA SOURCES

Psychostimulants

Our county-level data on psychostimulant distribution comes from the Drug Enforcement Administration's

Table 2. Characteristics of counties with high and low per capita use of psychostimulants, DEA's ARCOS database differences in means for all U.S. counties for methylphenidate and amphetamine combined (2000)

Variable	Mean for all U.S. counties (standard error)	Low consumption <i>n</i> = 1515 counties (standard error)	High consumption <i>n</i> = 1515 counties (standard error)
Socio-demographic and economic characteristics			
Psychostimulant distribution rate (g/per 100 000k) [†]	3359 g** (37.31)	1796 g** (20.72)	4923 g** (43.50)
Total population [†]	89 327 (52.22)	65 680 (86.25)	114 000 (59.34)
Per capita income [†]	\$21 397 (130.8)	\$20 118 (159.2)	\$22 760 (204.6)
Unemployment rate [†]	4.9% (0.049)	5.4% (0.079)	4.4% (0.055)
% of population with some form of health insurance [†]	82.9% (0.10)	82.1% (0.137)	83.7% (0.131)
White population as percentage of total population*	81.9% (0.290)	81.1% (0.455)	82.6% (0.362)
Black population as percentage of total population	12.7% (0.280)	13.2% (0.438)	12.4% (0.352)
Asian population as percentage of the total population	4.5% (0.280)	5.5% (0.441)	3.4% (0.358)
State has schedule II Rx monitoring program [†]	30.7% (0.008)	34.3% (0.012)	27.0% (0.012)
Children/adolescents as a % of the population [†]	26.0% (0.061)	26.2% (0.10)	25.8% (0.075)
Educational characteristics			
Students-to-teacher ratio [†]	14.6 (0.048)	14.3 (0.070)	14.8 (0.066)
Private students-to-public students ratio [†]	0.058 (0.001)	0.043 (0.0018)	0.069 (0.002)
Private coed-to-private non-coed students [†]	58.0 (8.609)	38.0 (8.092)	68.3 (12.33)
Private sectarian-to-private non-sectarian students [†]	23.8 (2.755)	14.5 (2.072)	28.5 (3.990)
Health system characteristics			
HMO penetration (% of individuals enrolled in HMOs) [†]	12.1% (0.248)	10.2% (0.322)	14.1% (0.374)
Number of HMOs operating in a county [†]	3.9 (0.069)	3.4 (0.092)	4.5 (0.101)
MDs/per 100 000 individuals [†]	112 (2.690)	86 (3.50)	139 (4.00)
Child psychiatrists as percentage of total MDs [†]	0.4% (0.026)	0.3% (0.038)	0.5% (0.036)
Psychiatrists as percentage of total MDs [†]	3.0% (0.099)	2.5% (0.149)	3.4% (0.119)
GPs, FPs as percentage of total MDs [†]	41.1% (0.515)	47.7% (0.757)	34.7% (0.662)
Pediatricians as percentage of total MDs [†]	5.5% (0.129)	4.9% (0.200)	6.1% (0.166)
Neurologists as percentage of total MDs [†]	0.9% (0.046)	0.7% (0.085)	1.1% (0.039)
Any psychiatrists present? [†]	0.544 (0.09)	0.684 (0.012)	0.401 (0.013)
Any child psychiatrists present? [†]	0.777 (0.007)	0.882 (0.008)	0.671 (0.012)
Any neurologists present? [†]	0.691 (0.008)	0.820 (0.009)	0.559 (0.012)
Any pediatricians present? [†]	0.434 (0.009)	0.554 (0.013)	0.307 (0.012)

(Continues)

Table 2. (Continued)

Variable	Mean for all U.S. counties (standard error)	Low consumption <i>n</i> = 1515 counties (standard error)	High consumption <i>n</i> = 1515 counties (standard error)
Female MDs-to-male MDs ratio	0.210 (0.004)	0.210 (0.006)	0.210 (0.004)
Younger MDs(<55)-to-senior MDs(>55) ratio [†]	1.8 (0.023)	1.6 (0.035)	1.9 (0.030)

Indicates that the variable's 'difference in means' is statistically significant at the 0.10 level.

*At the 0.05 level.

**g/per dosage unit.

[†]At the 0.01 level.

Automation of Reports and Consolidated Orders System (ARCOS) database, which monitors the flow of controlled substances from their point of manufacture through commercial distribution channels to points of sale or distribution at the dispensing/retail level: hospitals, retail pharmacies, practitioners and teaching institutions. Each year ARCOS reports more than 30 000 000 transactions.

Based on the DEA's description of the ARCOS database, distribution is highly related to consumption. The DEA's psychostimulant data that we analyzed includes the most commonly used drugs to treat ADHD: methylphenidate, amphetamines and dextroamphetamine sulfate.²⁰

For our area variation analysis in Table 2, we aggregated the DEA's 2000 ARCOS data on psychostimulant distribution from the five-digit zip code level to the county level (dependent variable) and then linked it with numerous county characteristics (independent variables) based on data from: area resource files (ARF), Census Estimates, InterStudy Publications on HMOs and the Department of Education's Common Core of Data. Approximately 70% of psychostimulants is prescribed for children and adolescents with attention deficit disorder (ADD).⁸ Thus, the majority of these drugs prescribed in any area are for persons younger than 19 years old and diagnosed with at least ADHD, as the diagnosis has high rates of comorbidity.⁹

In order to test the reliability and validity of the DEA's ARCOS data for measuring the area variability in psychostimulant use (not just distribution), we cross-checked it with auxiliary datasets that measure actual prescription amounts at the state level. For example, for the state of California we obtained prescription data from the state's Schedule II prescription drug monitoring program (run by the state's Department of Justice),¹⁰ which records information on every Schedule II prescription written within each of the state's 3-digit zip codes. The prescription claims data

from California's monitoring program lists the number of pills dispensed in each 3-digit zip code, and their various strengths, for a limited number of drugs containing methylphenidate (e.g. Concerta, Ritalin, methylphenidate hydrochloride etc.). Using this data, we classified each of the 57 3-digit zip codes in California as either 'high' (28 areas) or 'low' (29 areas) users of psychostimulants, depending on whether they were above or below the median level of consumption for these drugs in the state. We then took the DEA's ARCOS data for methylphenidate distribution to 5-digit zip codes in CA and aggregated it up to 3-digit level to match California's data.

Using the DEA's data, we again classified each of the 57 3-digit zip codes as either 'high' (again 28 areas) or 'low' (29 areas) use areas based on the median value. Of the 28 areas classified as 'high' use by data from California's prescription monitoring program, 27 were also correctly classified as 'high' use by the DEA data, while only 1 area was incorrectly classified as low use. Similarly, of the 29 areas classified by California's data as 'low' use areas, 28 were also classified as 'low' use by DEA, while one area was incorrectly classified as high use. Thus, the sensitivity and specificity of using DEA data is 96.4 and 96.6%, respectively. Because the incidence of high/low counties (using median values) is 0.5, this implies that the probability of false positives is only 0.0346 (or 3.5%). With this extremely low probability of false positives, we can safely use DEA data on distribution of psychostimulant drugs to classify all U.S. counties as having either 'high' or 'low' per-capita rates of psychostimulant consumption.

RESULTS

The county-level analysis of per-capita psychostimulant use reveals significant differences and similarities between 'low' and 'high' use areas. Counties with

higher per-capita consumption rates have, on average, greater population density, higher per capita income, lower unemployment rates, a greater proportion of their population with health insurance coverage, greater HMO penetration, more HMOs operating in each county and more physicians (including specialists) per capita. They do not, however, have more children as a proportion of the population.

Expressed as a percentage of total physicians, counties above the median rate of psychostimulant consumption have a greater percentage of pediatricians, psychiatrists, child psychiatrists and neurologists, but a lower percentage of general and family practitioners. While there appears to be no significant difference in the male-to-female ratio among physicians, counties with higher consumption rates have a higher young-to-old physician ratio.

There are also differences in the educational environment across the two sets of counties. Counties with higher rates of psychostimulant use have slightly higher students-to-teacher ratios in public schools, but significantly higher ratios of private school students-to-public schools students, private coed-to-private non-coed students and private sectarian (religious)-to-private non-sectarian students.

DISCUSSION

In general, the results in Table 2 suggest a positive relationship between county affluence and higher per-capita use of psychostimulant drugs. For instance areas with 'high' use of psychostimulants have, on average, about \$2650 greater per-capita income and a 1% lower unemployment rate than areas with 'low' use. These findings are consistent with the findings of a recent study on the geographic variation in the prevalence of psychostimulant medication use among children.²⁰ They also suggest a positive relationship between higher use of the drugs and areas with greater population density and greater access to medical care (more physicians, more specialists and higher levels of health insurance). To receive psychostimulants, one needs access to a prescribing physician. So it is not surprising that areas with greater affluence and health system resources also have higher use of psychostimulants.

What is surprising in our findings is the difference in the composition of medical professionals between areas of 'high' and 'low' psychostimulant use. While greater numbers of physicians per capita would be correlated with affluence, the areas with 'high' use of psychostimulants also have a higher ratio of younger-to-older physicians: areas with 'high' use have 1.9 younger (<55 years old) physicians for every 1 older

(>55 years old) physician, whereas in areas with 'low' use the ratio is only 1.6 to 1. Two plausible explanations for this difference are that: (1) younger physicians are systematically self-locating in more affluent areas (which are correlated with high use of psychostimulants), and/or (2) that the medical training of younger physicians is inherently different and that, on average, they are more likely than their older colleagues to prescribe psychostimulants. While the first explanation is certainly plausible, we believe that in recent years—with ADHD being more widely recognized and accepted as a legitimate mental disorder—the second explanation probably explains a greater proportion of the variance. If this is the case, then it behooves us to further investigate if older physicians are more likely to be 'under' prescribing or if younger physicians are more likely to be 'over' prescribing. Similar differences can be seen across various medical specialties. Areas with 'high' use have a greater percentage of psychiatrists, child psychiatrists and pediatricians as a percentage of total physicians in the area.

Another important difference between areas with 'high' and 'low' use is both the number of HMOs and total HMO penetration (where penetration is measured as the percentage of the population enrolled in HMOs). Given the cost-minimizing objectives of HMOs, one would expect the consumption of psychostimulants to be lower in areas with higher HMO penetration. However, the results show that areas with 'high' consumption have greater HMO penetration. Once again, this could be simply a manifestation of spurious correlations with other variables (e.g. HMOs systematically entering more affluent markets). Yet this finding is also consistent with the cost minimization objectives if we consider that the alternative treatment—counseling and/or psychiatric therapy that focuses on cognitive and behavioral adjustments—is far more costly. Thus, HMOs could be systematically substituting cheaper drug therapy for the more expensive counseling options. (Note that HMOs advocate the use of one product over another via the formularies; however, this does not affect our finding because our results are not for a specific psychostimulant drug such as Ritalin or Adderall, but rather for all psychostimulants combined.)

In addition, we find that 'high' use counties have, on average, slightly fewer children as a proportion of the population. One would expect that areas with a higher proportion of children would have higher per-capita rates of psychostimulant use, given that ADHD is primarily diagnosed among children. A possible explanation for this counterintuitive result is provided

by a recent research finding that children in families with four or more children were 26% less likely than children in smaller families to consume a psychostimulant medication.²⁰ In other words, areas that have smaller families are more likely to have fewer children as a proportion of the total population, but the children are more likely to be using these drugs.

Finally, we find that areas with 'high' use have a slightly higher students-to-teacher ratio than areas with 'low' use. This finding is in sharp contrast to the affluence explanation given above, since more affluent areas typically have lower students-to-teacher ratios. Again, there are two competing explanations for this finding: (1) areas where students-to-teacher ratios are higher place a greater burden on teachers; hence, misbehavior could lead to more children being sent to the school psychologist for screening, resulting in higher detection rates of ADHD and ultimately of psychostimulant consumption; (2) that the observed correlation is just that, a (spurious) correlation—controlling for other factors will make this finding insignificant. Given that population is greater in 'high' use counties, we tend to favor the second explanation.

LIMITATIONS

One limitation of this study, as previously mentioned, is that the ARCOS data measures distribution of drugs (not actual consumption). For example, the shipment may be sent to one location, but used in another. Our use of other auxiliary datasets, however, which do measure psychostimulant consumption at the state level, confirms that the DEA data is a sufficiently accurate measure of the variability in the use of these drugs (albeit not an ideal one).

A second limitation of this study, also previously mentioned, is that it is merely descriptive with bivariate correlates. Because we do not have price data that would show how much the price of these drugs varies geographically, we concluded that we could not perform a multivariate analysis without undue concern over potential omitted variable bias. In other words, we assume that the price of psychostimulant drugs has some measure of impact on its consumption. And not being able to account for it, therefore, casts doubt on the results of any multivariate analysis. In addition to our own efforts, we encourage others to try to obtain compatible price data that can be used to determine which factors (socio-demographic, educational or health system) are the most significant, when others are controlled, in predicting counties' per-capita psychostimulant use.

DIRECTIONS FOR FUTURE RESEARCH

While there is significant variation in the use of psychostimulant drugs across the country, there is also considerable variation in other variables as well. An important and necessary direction for future research is to test if any of these covariates are causally related to consumption rates and, if so, the direction and strength of causality should be estimated. A starting point is to specify a structural model for the equilibrium quantity and price of the psychostimulant drugs in each market and then test hypotheses regarding the effects of these covariates on the consumption rates. The difficulty, however, lies in obtaining price data on psychostimulant drugs at the county level to match the DEA's quantity data. As previously explained, this is why we restricted our study to a bivariate analysis.

Ultimately, differences in psychostimulant consumption rates across the country could arise for three basic reasons: a greater proportion of people are diagnosed with ADHD in some areas than in others;^{14,17–20} a greater proportion of people are prescribed psychostimulants in some areas than in others, even though the same proportion of people may be diagnosed with ADHD;⁸ and the same proportion of people are diagnosed with ADHD and prescribed psychostimulants, but patients in some areas are systematically given larger dosages than in other areas.^{14,16}

Given these reasons and our preliminary findings, we have identified two areas ripe for additional research. First, market level, multivariate analyses (which control for price differentials) are needed to tease out how—and to what extent—economic, demographic, educational, regulatory and health system factors affect regional rates of psychostimulant consumption. Second, researchers should examine physicians' practice styles in diagnosing and treating ADHD to see how much they vary, and whether they vary systematically by patients' and/or physicians' personal characteristics (e.g. age, race, gender, ethnicity, type of health insurance, professional experience, type of practice—group/solo etc.).

ACKNOWLEDGEMENTS

We wish to thank Doug Schwalm, Ph.D., James Bosco, Ph.D., Patricia Pastor, Ph.D., and Jennifer Mayes for their helpful critiques of this paper at various stages of its development. We also want to express our gratitude to the Nicholas C. Petris Center on Healthcare Markets & Consumer Welfare and the Center for Child & Youth Policy at the University of California, Berkeley for their financial support of our research.

REFERENCES

- Cantwell D. Attention deficit disorder: a review of the past 10 years. *J Am Acad Child Adolescent Psychiatry* 1996; **35**: 978–987.
- LeFever G, Dawson K, Morrow A. The extent of drug therapy for attention deficit-hyperactivity disorder among children in public schools. *Am J Pub Health* 1999; **89**: 1359–1364.
- National Institutes of Health (NIH). *Consensus Statement Online, Diagnosis and Treatment of Attention Deficit Hyperactivity Disorder, Consensus Development Conference Statement* [online]. Available: http://odp.od.nih.gov/consensus/cons/110/110_statement.htm [16–18 November 1998].
- Guevara J, Lozano P, Wickizer T, Mell L, Gephart H. Utilization and cost of health care services for children with attention-deficit/hyperactivity disorder. *J Pediatrics* 2001; **108**: 71–78.
- Jensen P. Commentary: the NIH ADHD consensus statement: win, lose, or draw? *J Am Acad Child Adolescent Psychiatry* 2000; **39**: 194–197.
- Zito J, Safer D, dosReis S, Magder L, Gardner J, Zarin D. Psychotherapeutic medication patterns for youths with attention-deficit/hyperactivity disorder. *Arch Pediatrics Adolesc Med* 1999; **153**: 1257–1263.
- Sloan M, Jensen P, Kettle L. Assessing services for children with ADHD: gaps and opportunities. *J Attention Disorders* 1999; **3**: 13–29.
- Safer D, Zito J, Fine E. Increased methylphenidate usage for attention deficit disorder in the 1990s. *Pediatrics* 1996; **98**: 1084–1088.
- Cantwell D, Baker L. Association between attention deficit hyperactivity disorder and learning disorders. *J Learn Disabil* 1991; **24**: 88–94.
- State prescription monitoring programs track the quantity of the drug and the identity of the physician and patient for every individual psychostimulant prescription filled by a pharmacy. Control begins with the establishment of production limits (quotas) for Schedule I & II substances. Schedule I substances have the highest potential for abuse and addiction and have no accepted medical use in the United States. Schedule II includes such drugs as methadone, meperidine (Demerol), cocaine, oxycodone (Percodan) and methylphenidate (Ritalin). Because of their high abuse potential, the use of Schedule II drugs is the most restricted and greater controls are placed on them.
- Zito J, Safer D, dosReis S, Gardner J, Bolese M, Lynch F. Trends in the prescribing of psychotropic medications to preschoolers. *JAMA* 2000; **283**: 1025–1032.
- Searight H, Burke J, Rottnek F. Adult attention deficit hyperactivity disorder (ADHD): evaluation and treatment in family medicine. *Am Family Physician* 2000; **62**: 2077–2086, 2091–2092.
- Rappley M, Gardiner J, Jetton J, Houang R. The use of methylphenidate in Michigan. *Arch Gen Psychiatr* 1995; **149**: 675–679.
- Spanos B. Quotas, ARCOS, UN Report and Statistics. Conference Report: stimulant Use in the Treatment of ADHD. Washington, D.C.: Drug Enforcement Administration, Department of Justice; 1996.
- Glass C, Wegar K. Teacher perceptions of the incidence and management of attention deficit hyperactivity disorder. *Educacion* 2000; **121**: 412–420.
- Forness S, Kavale K. ADHD and a return to the medical model of special education. *Educ Treatment Children* 2001; **24**: 224–247.
- Safer D, Krager J. Effect of a media blitz and a threatened lawsuit on stimulant treatment. *JAMA* 1992; **268**: 1004–1007.
- Zito J, Safer D, dosReis S, Magder L, Gardner J, Zarin D. Psychotherapeutic medication patterns for youths with attention-deficit/hyperactivity disorder. *Arch Pediatrics Adolesc Med* 1999; **153**: 1257–1263.
- Zito J, Safer D, dosReis S, Magder L, Riddle M. Methylphenidate patterns among medicaid youths. *Psychopharmacol Bull* 1997; **30**: 143–147.
- Cox E, Motheral B, Henderson R, Mager D. Geographic variation in the prevalence of stimulant medication use among children 5 to 14 years old: results from a commercially insured sample. *J Pediatrics* 2003; **111**: 237–243.
- Bird HR, Canino G, Rubio-Stipec M, et al. Estimates of the prevalence of childhood maladjustment in a community survey in Puerto Rico: the use of combined measures [published correction appears in *Archives of General Psychiatry*. 1994; **51**: 429]. *Arch Gen Psychiatry* 1988; **45**: 1120–1126.
- Cohen P, Cohen J, Kasen S, et al. An epidemiological study of disorders in late childhood and adolescence, I: age- and gender-specific prevalence. *J Child Psychol Psychiatry* 1993; **34**: 851–867.
- Costello EJ, Costello AJ, Edelbrock C, et al. Psychiatric disorders in pediatric primary care: prevalence and risk factors. *Arch Gen Psychiatry* 1988; **45**: 1107–1116.
- Esser G, Schmidt MH, Woerner W. Epidemiology and course of psychiatric disorders in school-aged children: results of a longitudinal study. *J Child Psychol Psychiatry* 1990; **31**: 243–263.
- Offord DR, Boyle MH, Szatmari P, et al. Ontario Child Health Study, II: six month prevalence of disorder and rates of service utilization. *Arch Gen Psychiatry* 1987; **44**: 832–836.
- Reid R, Maag JW, Vasa SF, Wright G. Who are the children with attention deficit-hyperactivity disorder? A school-based survey. *J Special Educ* 1994; **28**: 117–137.
- Szatmari P, Offord DR, Boyle MH. Ontario Child Health Study: prevalence of attention deficit disorder with hyperactivity. *J Child Psychol Psychiatry* 1989; **30**: 219–230.
- Velez CN, Johnson J, Cohen P. A longitudinal analysis of selected risk factors for childhood psychopathology. *J Am Acad Child Adolesc Psychiatry* 1989; **28**: 861–864.
- IMS Health. Retail and Provider Perspective Audit, 1990–2000; and National Disease and Therapeutic Index, 1980–2000. Plymouth Meeting, Penn. 2001.
- Swanson J, Lerner M, Williams L. More frequent diagnosis of attention deficit-hyperactivity disorder. *New Engl J Med* 1995; **333**: 944.
- State prescription monitoring programs track the quantity of the drug and the identity of the physician and patient for every individual psychostimulant prescription filled by a pharmacy. Control begins with the establishment of production limits (quotas) for Schedule I & II substances. Schedule I substances have the highest potential for abuse and addiction and have no accepted medical use in the United States. Schedule II includes such drugs as methadone, meperidine (Demerol), cocaine, oxycodone (Percodan) and methylphenidate (Ritalin). Because of their high abuse potential, the use of Schedule II drugs is the most restricted and greater controls are placed on them. Several states have instituted multiple copy prescription programs or electronic tracking systems, which can be used alone or with the single-serialized form. Both systems allow states to monitor these drugs. A multiple copy prescription monitoring program operates as follows: (1) The prescriber writes a prescription for a Schedule II (and in a few states, certain

VARIATION IN PSYCHOSTIMULANT USE

Schedule III and IV) controlled substance on a state issued, preprinted, serialized duplicate or triplicate form; (2) The prescriber writes and the dispenser maintains file copies of the prescription for a period of two to five years (for triplicate programs). Duplicate prescription programs do not require the prescriber to maintain copies; (3) The dispenser forwards a copy of the prescription to the mandated state authority; With an electronic data transmission system, however: (1) The prescriber writes an original prescription for a Schedule II (and in some states, Schedules III, IV and V) controlled substance on a prescription form; (2) The dispenser maintains the original prescription for a period of two to five years; (3) The dispenser

transmits the prescription information either electronically (via modem, disk, tape, black box) or by universal claim form to the mandated state authority. This system allows prescription information to be submitted electronically. In most states, if the dispenser lacks the requisite computer equipment and/or fills less than 20-25 Schedule II prescriptions per month, information is submitted on a Universal Claim Form.

32. U.S. General Accounting Office (GAO). Prescription Drugs: State Monitoring Programs Provide Useful Tool To Reduce Diversion. Report to the Subcommittee on Oversight and Investigations, Committee on Energy and Commerce, House of Representatives, GAO-02-634. Washington, D.C.; May 2002.